
# **CONCEPT: SOUTHERN BLOTTING**

## **Radioactive Probes**

<ul> <li>After cloning a gene, th</li> </ul>	e DNA can be used as a <mark>probe</mark> to detect the	sequence in an unknown DNA sample.
;	radioactively labeled molecules that are visualized using ra	adioactive detection.
□ DNA probes a	arestranded DNA molecules complementar	ry to a <i>specific</i> sequence of interest.

**EXAMPLE:** Radioactive probes are used to identify DNA that is complementary to it in different samples.



**PRACTICE:** A southern blotting technique is used to detect a specific \_\_\_\_\_\_ sequence from a blood or tissue sample.

a) RNA

c) Protein

b) DNA

d) Lipid

**PRACTICE:** A \_\_\_\_\_ is a single-stranded DNA molecule used in hybridization reactions to detect the presence of a particular gene in an assortment of DNA fragments.

(Hybridization Reactions: reactions that combine two complementary single-stranded DNA molecules)

- a) Plasmid.
- b) Vector.
- c) Probe.
- d) Blot.
- e) Polymerase.

#### **CONCEPT: SOUTHERN BLOTTING**

<ul> <li>Blotting: technique used to rapidly detect a specific DNA sequence using DNA pro</li> </ul>			
	□ Northern Blotting detects	sequences & Western Blotting detects specific	

## Steps of Southern Blotting

1			
1	4	Unknown DNA sample is fragmented & separated by	y using gel electrophoresis.
L	-	OHKHOWH DINA Samble is hadillented & Sebarated by	v usiilu dei electroprioresis.

2 DNA on the gel is *denatured* to \_\_\_\_\_\_-stranded DNA (ssDNA) by incubating it with *denaturing buffer*.


□ **Filter paper** used to "*blot*" the gel & absorb the *denaturing buffer* as it migrates through to a paper towel stack.

3 ssDNA is \_\_\_\_\_ to the nitrocellulose filter paper when denaturing buffer is absorbed/migrated.

4 Filter paper is removed & incubated with the radioactive probe \_\_\_\_\_\_\_to the sequence of interest.

5 Radioactive filter paper is analyzed & visible bands are ones that \_\_\_\_\_\_ to the DNA probe in step 4.

**EXAMPLE:** Analyze the Southern Blot below and decide which samples contain the DNA sequence of interest.



**PRACTICE:** A geneticist wants to see if her patient has Gene X. The geneticist takes a blood sample from her patient and prepares a southern blot. How will the geneticist know if her patient possesses Gene X?

- a) A radioactive probe, which is complementary to Gene X, will bind to the membrane.
- b) A radioactive probe, which has the same sequence as Gene X, will bind to the membrane.
- c) A radioactive probe will be absent from the membrane.
- d) A southern blot cannot confirm if the patient possesses Gene X.

### **CONCEPT: SOUTHERN BLOTTING**

**PRACTICE:** Place the following steps of Southern Blotting in the correct order.

- a) \_\_\_\_\_: Filter paper is incubated with the labeled DNA probe which anneals to the ssDNA fragments.
- b) \_\_\_\_\_: Analyze gel to determine the presence of the DNA sequence of interest.
- c) \_\_\_\_\_: Separate DNA fragments by size using gel electrophoresis.
- d) \_\_\_\_\_: Denature DNA by soaking gel in a basic solution.
- e) \_\_\_\_\_: Fragment unknown DNA sample(s) using restriction enzymes.

PRACTICE: By analyzing the Southern Blot results below, which of the samples contains the gene of interest?

- a) A only.
- b) A & C.
- c) B & D.
- d) A, C, & E.
- e) E only.
- f) None.

