


## CONCEPT: EUKARYOTIC POST-TRANSLATIONAL REGULATION

- Eukaryotes regulate expression at the post-\_\_\_\_\_ by controlling activity of the expressed protein.
  - Recall: *Post-translational modifications (PTMs)* are covalent modifications to proteins \_\_\_\_\_ translation.
- PTMs can activate/inactivate a protein or “\_\_\_\_\_” the protein for degradation by **Proteases**.
  - \_\_\_\_\_: enzymes that degrade proteins by breaking *polypeptide bonds* making single amino acids.

**EXAMPLE:** Protein activity can be controlled by post-translational modifications or degradation by proteases.



**PRACTICE:** Protein degradation is one strategy to control gene expression and is considered \_\_\_\_\_.

- Transcriptional control.
- Post-transcriptional control.
- Translation initiation control.
- Post-translational control.
- Chromatin remodeling.

**PRACTICE:** Post-translational modifications of proteins can affect which of the following?


- Protein function.
- Protein location within the cell.
- Protein activation or inactivation.
- Protein degradation.
- All of the above.

## CONCEPT: EUKARYOTIC POST-TRANSLATIONAL REGULATION

### Protein Ubiquitination

- Eukaryotes need a way to \_\_\_\_\_ which proteins in a cell are no longer needed & can be removed.
- Cells utilize PTMs to “tag” specific proteins in a cell to be \_\_\_\_\_ by cellular *proteases*.
  - \_\_\_\_\_: small peptide used by Eukaryotic cells to mark proteins for degradation.
  - **Ubiquitin** \_\_\_\_\_: cellular enzyme that adds the *ubiquitin* peptide to the target protein.

**EXAMPLE:** Ubiquitin ligase adds a ubiquitin peptide to a mis-folded or non-functioning protein.



**PRACTICE:** A hormone signal reaches a cell and causes the cell to produce a large quantity of Protein X. After some time, the hormone signal disappears and the cell no longer needs a large quantity of Protein X. How will the cell remove the excess protein?

- The repressor protein for the Protein X gene will stop the transcription of the gene.
- The excess Protein X will be tagged with ubiquitin proteins and degraded over time.
- The Protein X mRNA will be bound by a microRNA blocking its translation.
- Over time the excess Protein X will diffuse out of the cell.