

TOPIC: AVERAGE VALUE OF A FUNCTION

Average Value of a Function

◆ Recall: Taking the limit as $n \rightarrow \infty$ of a Riemann sum gives the *true* area under the curve: the definite integral.

► We can use a definite integral to find the average value of a function $f(x)$ from $[a, b]$.

New

Average Value of a Function

$$\text{Avg value} \approx \frac{f(x_1^*) + f(x_2^*) + \cdots + f(x_n^*)}{n}$$

$$\text{Avg value} \approx \frac{1}{n} \sum_{k=1}^n f(x_k^*)$$

$$\text{Avg value} \approx \frac{1}{n} \sum_{k=1}^n f(x_k^*)$$

$$\text{Avg value} \approx \frac{1}{n} \sum_{k=1}^n f(x_k^*)$$

$$\text{Avg value} = \frac{1}{b-a} \cdot \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k^*) \Delta x$$

$$f_{\text{avg}} = \frac{1}{b-a} \int_a^b f(x) dx$$

Recall

$$\Delta x = \frac{b-a}{n}$$

Recall

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(x_k^*) \Delta x$$

EXAMPLE

Find the average value of $f(x) = x + 2$ on $[0, 4]$.

TOPIC: AVERAGE VALUE OF A FUNCTION

PRACTICE

Find the average value of the function on the interval $[2,5]$.

$$F(x) = x^3 - \frac{2}{\sqrt{x}}$$

PRACTICE

Find the average value of the function on the interval $[4,9]$.

$$F(x) = \sqrt{x} + 6$$

TOPIC: AVERAGE VALUE OF A FUNCTION

PRACTICE

Find the average value of the function on the interval $[0,1]$.

$$G(x) = \frac{2}{x^2 + 1}$$